CONTENTS

	Page
Relation of Carbon Bisulphid to Soil Organisms and Plant Growth. E. B. Fred	1
Climatic Conditions as Related to Cercospora beticola. Venus W. Pool and M. B. McKay	21
Soluble Nonprotein Nitrogen of Soil. R. S. Potter and R. S. Snyder	61
Oviposition of Megastigmus spermotrophus in the Seed of Douglas Fir. J. M. Miller	65
Citrus Canker. Frederick A. Wolf	69
Determination of Stearic Acid in Butter Fat. E. B. Holland, J. C. Reed, and J. P. Buckley, Jr	101
Life History and Habits of Two New Nematodes Parasitic on Insects. J. H. Merrill and A. L. Ford	115
Insect Injury to Cotton Seedlings. B. R. Coad and R. W. Howe	129
A Sex-Limited Color in Ayrshire Cattle. Edward N. Wentworth	141
Watermelon Stem-End Rot. F. C. Meier	149
Effect of Pasteurization on Mold Spores. Charles Thom and S. Henry Ayers	153
Effect of Water in the Ration on the Composition of Milk. W. F. Turner, W. H. Shaw, R. P. Norton, and P. A. Wright	167
Crowngall Studies, Showing Changes in Plant Structures Due to a Changed Stimulus. Erwin F. Smith	179
Effect of Certain Species of Fusarium on the Composition of the Potato Tuber. Lon A. Hawkins	183
Hyperaspis binotata, a Predatory Enemy of the Terrapin Scale. F. L. Simanton	197
Test of Three Large-Sized Reinforced-Concrete Slabs under Concentrated Loading. A. T. Goldbeck and E. B. Smith	205
Occurrence of Sterile Spikelets in Wheat. A. E. Grantham and Frazier Groff	235
Effect of Cold-Storage Temperatures upon the Pupæ of the Mediterranean Fruit Fly. E. A. Back and C. E. Pemberton	251
Effect of Climatic Factors on the Hydrocyanic-Acid Content of Sorghum. J. J. Willaman and R. M. West	261
Egg and Manner of Oviposition of Lyctus planicollis. Thomas E. Snyder	273
Hypoderma deformans, an Undescribed Needle Fungus of Western Yellow Pine. James R. Weir	277

Page

Ornix germinatella, the Unspotted Tentiform Leaf Miner of Apple. L. HASEMAN 289
A Western Fieldrot of the Irish Potato Caused by Fusarium radicicola. O. A. PRATT 297
Comparative Study of the Root Systems and Leaf Areas of Corn and the Sorghums. E. C. MILLER 311
Production of Clear and Sterilized Anti-Hog-Cholera Serum. M. DORSET and R. R. HENLEY 333
Silver-Scurf of the Irish Potato Caused by Spondylocladium atrovirens. EUGENE S. SCHULTZ 339
Woolly Pear Aphis. A. C. BAKER and W. M. DAVIDSON 351
Pathological Histology of Strawberries Affected by Species of Botrytis and Rhizopus. NEIL E. STEVENS 361
Life Histories and Methods of Rearing Hessian-Fly Parasites. C. M. PACKARD 367
Effect of Röntgen Rays on the Tobacco, or Cigarette, Beetle and the Results of Experiments with a New Form of Röntgen Tube. G. A. RUNNER 383
Stimulating Influence of Arsenic upon the Nitrogen-Fixing Organisms of the Soil. J. E. GREAVES 389
Transmission and Control of Bacterial Wilt of Cucurbits. FREDERICK V. RAND and ELLA M. A. ENLOWS 417
Correlated Characters in Maize Breeding. G. N. COLLINS 435
Comparative Study of the Amount of Food Eaten by Parasitized and Nonparasitized Larvæ of Cirphis unipuncta. DANIEL G. TOWER 455
Aleyrodidae, or White Flies Attacking the Orange, with Descriptions of Three New Species of Economic Importance. A. L. QUAINTANCE and A. C. BAKER 459
Relative Water Requirement of Corn and the Sorghums. EDWIN C. MILLER 473
Availability of Mineral Phosphates for Plant Nutrition. W. L. BURLISON 485
California Green Lacewing Fly. V. L. WILDERMUTH 515
Rape as Material for Silage. A. R. LAMB and JOHN M. EVVARD 527
Effect of Autolysis upon Muscle Creatin. RALPH HOAGLAND and C. N. MCBRYDE 535
Storage Rots of Economic Aroids. L. L. HARTER 549
Experiments with Clean Seed Potatoes on New Land in Southern Idaho. O. A. PRATT 573
Digestibility of Very Young Veal. C. F. LANGWORTHY and A. D. HOLMES 577
Influence of Calcium and Magnesium Compounds on Plant Growth. F. A. WYATT 589

Page

Larval Characters and Distribution of Two Species of Diatraea. T. E. HOLLOWAY 621
The Disease of Potatoes Known as "Leak." LON A. HAWKINS.. 627
Digestibility of Hard Palates of Cattle. C. F. LANGWORTHY and A. D. HOLMES 641
Some Properties of the Virus of the Mosaic Disease of Tobacco. H. A. ALLARD 649
Life Cycles of the Bacteria. F. LÖHNIS and N. R. SMITH 675
A Respiration Calorimeter, Partly Automatic, for the Study of Metabolic Activity of Small Magnitude. C. F. LANGWORTHY and R. D. MILNER 703
Mottle-Leaf of Citrus Trees in Relation to Soil Conditions. LYMAN J. BRIGGS, C. A. JENSEN, and J. W. MCLANE 721
Vegetative Succession Under Irrigation. J. FRANCIS MACBRIDE.. 741
Agricultural Value of Impermeable Seeds. GEORGE T. HARRINGTON 761
Mendelism of Short Ears in Sheep. E. G. RITZMAN 797
Life-History Studies of Cirphis Unipuncta, the True Army Worm. JOHN J. DAVIS and A. F. SATTERTHWAIT 799
Infection of Timothy by Puccinia graminis. E. C. STAKMAN and F. J. PIEMEISEL 813
Control of the Powdery Dryrot of Western Potatoes Caused by Fusarium trichothecioides. O. A. PRATT 817
Use of the Moisture Equivalent for the Indirect Determination of the Hygroscopic Coefficient. FREDERICK J. ALWAY and JOUETTE C. RUSSEL 833
Thersilochus conotracheli, a Parasite of the Plum Curculio. R. A. CUSHMAN 847
Effect on Plant Growth of Sodium Salts in the Soil. FRANK B. HEADLEY, E. W. CURTIS, and C. S. SCOFIELD 857
Observations on the Life History of the Army Cutworm, Chorizagrotis auxiliaris. R. A. COOLEY 871
Aphidoletes meridionalis, an Important Dipterous Enemy of Aphids. JOHN J. DAVIS 883
Influence of Barnyard Manure and Water upon the Bacterial Activities of the Soil. J. E. GREAVES and E. G. CARTER 889
Progressive Oxidation of Cold-Storage Butter. D. C. DYER 927
Bacteriological Studies of a Soil Subjected to Different Systems of Cropping for Twenty-five Years. P. L. GAINEY and W. M. GIBBS 953
Studies on the Physiology of Reproduction in the Domestic Fowl.—XV. Dwarf Eggs. RAYMOND PEARL and MAYNIE R. CURTIS.. 977
a-Crotonic Acid, a Soil Constituent. E. H. WALTERS and LOUIS E. WISE 1043
Index 1047

ERRATA.

Page 151, line 25, "Sphaeropsideae" should read "Sphaerioidaceae."

Page 156, "*Penicillium camemberti*, var. *rogri*" should read "*Penicillium camemberti*, var. *rogeri*."

Page 296, Pl. XXXIII, figs. 3 to 15. The magnification of the illustrations should be half that stated in the legend.

Page 303, line 17, "Plate XXXVI, figures 1 to 4" should read "Plate XXXVII, figures 1 to 4" and "In figure 4, Plate XXXVI" should read "In figure 4, Plate XXXVII."

Page 318, Table IV, under head "General remarks," "rooting" should read "shooting."

Page 337, Table II, 4th column, "Phenolized defibrinated blood 3895 (unwashed)" should read "Phenolized defibrinated blood 3895."

Page 377, last line, 2d paragraph, "winged" should read "wingless."

Page 384, Table I, 6th column, "Current (milliampere minutes)" should read "Current (milliamperes)."

Page 388, line 13 from bottom, omit "with humidity at 57."

Page 419, line 25, "The twelve-spotted (or squash) lady beetle" should read "The squash lady beetle."

Page 419, line 28, "(*Crepidodera cucumeris*)" should read "(*Epitrix cucumeris*)."

Page 459, lines 2 and 24, omit "Three."

Page 471, line 4, "*Aleurodes mori* Ckll." should read "*Aleurodes mori*, var. *arizonensis* Ckll."

Page 762, Table I, first column, "*Medicago arbica*" should read "*Medicago arabica*."

Page 791, Table XIV, 1st column, "(p. 23)" should read "(p. 783)."

Page 865, legend under figure 5, "The solid black line, etc.," should read "The hatched line."

Page 866, legend under figure 6, omit sentences 2 and 3.

Page 881, line 6 from bottom, "April" should read "May."

ILLUSTRATIONS

PLATES

Relation of Carbon Bisulphid to Soil Organisms and Plant Growth

Page

Plate I. Plate cultures of soil organisms growing on agar: Fig. 1.—Colonies of organisms from untreated soil. Fig. 2.—Colonies from soil treated with 2 per cent of carbon bisulphid. Fig. 3.—Colonies from soil treated with 2 per cent of carbon bisulphid and evaporated. Fig. 4.—Colonies from soil treated with 2 per cent of carbon bisulphid, evaporated, and reinoculated with 5 per cent of soil from an untreated jar 20

Plate II. Fig. 1.—Effect of varying amounts of carbon bisulphid on mustard; *A*, *B*, soil untreated; *C*, *D*, soil treated with 0.5 per cent of carbon bisulphid; *E*, *F*, soil treated with 1 per cent of carbon bisulphid; *G*, *H*, soil treated with 2 per cent of carbon bisulphid. Fig. 2.—Effect of carbon bisulphid on clover in peat soil; *A*. *B*, soil untreated; *C*, *D*, soil treated with 2 per cent of carbon bisulphid. Fig. 3.—Effect of carbon bisulphid on buckwheat in sand cultures; *A*, *B*, soil untreated; *C*, *D*, soil treated with 2 per cent of carbon bisulphid 20

Climatic Conditions as Related to Cercospora beticola

Plate III. *Cercospora beticola:* Overwintering tests on the experimental field at Rocky Ford, Colo., during 1912–13: Sugar-beet leaves infected with *Cercospora beticola* (*1*) stored in soil in boxes, (*2*) buried in the ground at different depths from 1 to 8 inches, and (*3*) left exposed above the ground in a pile of hayed sugar-beet tops 60

Plate IV. Field stations for the collection of weather data at Rocky Ford, Colo., in 1913: Fig. 1.—Weather shelter, anemometer, and rain gauge at edge of sugar-beet field. Fig. 2.—Weather shelter among beet plants, showing hygrothermograph and cog psychrometer. Fig. 3.—Weather shelter of the local Weather Bureau station about 3 miles from sugar-beet field 60

Oviposition of Megastigmus spermotrophus in the Seed of Douglas Fir

Plate V. Oviposition of *Megastigmus spermotrophus* in the cones of Douglas fir: Fig. 1.—Type of cage in which the oviposition of *Megastigmus spermotrophus* was observed. Fig. 2, 3.—Female resting on cone with ovipositor inserted 68

Plate VI. Oviposition of *Megastigmus spermotrophus* in the cones of Douglas fir: Figs. 1, 2.—Two positions of female on surface of cone with ovipositor inserted. Fig. 3.—Female resting on cone with ovipositor inserted 68

Plate VII. Oviposition of *Megastigmus spermotrophus* in the cones of Douglas fir: Fig. 1.—Female in act of withdrawing ovipositor from cone. Fig. 2.—Section through a Douglas fir cone on which a female has been killed while in the act of ovipositing. Fig. 3.—A portion of same cone and dead female with ovipositor inserted 68

CITRUS CANKER

Page

PLATE VIII. Fig. 1.—Grapefruit leaf showing young Citrus cankers. Fig. 2.—Old Citrus canker on Satsuma leaves. Figs. 3, 4.—Seedling grapefruit branches affected with Citrus canker. Fig. 5.—Severe canker infection of branches of *Citrus trifoliata*..... 100

PLATE IX. Fig. 1.—View of lower side of leaves of seedling grapefruit artificially inoculated with *Pseudomonas citri*. Fig. 2.—Top view of plant shown in figure 1. Fig. 3.—Spongy white cankers on leaf and twig of seedling grapefruit produced by artificial inoculation. Fig. 4.—Citrus canker on Satsuma leaves resulting from artificial inoculation with *Pseudomonas citri*. Fig. 5.—Photomicrograph of section of young, open canker on grapefruit.. 100

PLATE X. Fig. 1.—Natural Citrus canker infection on leaves of *Citrus trifoliata*. Fig. 2.—Mature cankers on fruit of *Citrus decumana*. Fig. 3.—Canker on seedling grapefruit leaves, entrance having been effected through abrasions made by thorns. Fig. 4.—Young spongy cankers on fruit of *Citrus decumana*. Fig. 5.—*Phoma socia* on cellulose agar showing dissolution of cellulose. Fig. 6.—Mature cankerous areas on leaves of Duncan grapefruit.... 100

PLATE XI. Fig. 1.—Cankers on old grapefruit leaves which have enlarged during the second growing season. Fig. 2.—Citrus canker resulting from immersion of leaves in a bacterial suspension. Fig. 3.—Culture of *Phoma socia* showing pycnidial formation in concentric rings. Fig. 4.—Dilution poured plate of *Pseudomonas citri* on green-bean agar. Fig. 5.—Photomicrograph of pycnidium of *Phoma socia* taken in reflected sunlight. Fig. 6.—Photomicrograph of pycnidia of *Phoma socia* taken in diffuse light..... 100

INSECT INJURY TO COTTON SEEDLINGS

PLATE XII. Fig. 1.—Cutworm injury to cotton seedlings; produced in breeding cages. Figs. 2, 3.—Cutworm injury to cotton seedling..... 140

PLATE XIII. Fig. 1.—Cutworm injury to cotton seedling. Fig. 2.—Tussock larva feeding upon cotton leaf. Fig. 3.—Injury produced by a nearly full-grown tussock larva when confined in a screen cage containing potted cotton plants..... 140

PLATE XIV. Cotton leaves showing grasshopper injury..... 140

PLATE XV. Fig. 1.—Underside of cotton leaf showing grasshopper injury. Fig. 2.—Cotton leaf showing grasshopper injury..... 140

PLATE XVI. Fig. 1.—Injury to terminal bud of cotton by lepidopterous larva. Fig. 2.—Two cotton plants from laboratory garden with leaves removed.. 140

WATERMELON STEM-END ROT

PLATE XVII. Watermelons, showing the effect of inoculation with species of Diplodia: Fig. 1.—The upper four melons were held as checks; the lower five are melons nine days after having been inoculated with a culture of *Diplodia* sp. Fig. 2.—A watermelon nine days after having been inoculated with a culture of *Diplodia tubericola* E. and E..... 152

CROWNGALL STUDIES SHOWING CHANGES IN PLANT STRUCTURES DUE TO A CHANGED STIMULUS

PLATE XVIII. Teratoid crowngalls produced in *Pelargonium* spp..... 182

PLATE XIX. Teratoid crowngalls produced in castor-oil plant. Fig. A.—A red-stem variety. Leaves reflexed; axis distorted; and feeble shoots developing out of the axillary tumors. Fig. B.—A green-stem glaucous variety..... 182

Page

PLATE XX. Teratoid crowngalls produced in tobacco by inoculating *Bacterium tumefaciens*..... 182

PLATE XXI. The teratoid tumor strand of Plate XX, which gives rise during its course to more than 30 small tumors. Top.—Cross section of outer part of right side of stem of tobacco plant shown on Plate XX. *P*, outer edge of the phloem; *E*, epidermis; *T St*, tumor strand, which is bedded in the normal cortex of the stem. Bottom.—Longitudinal section from upper part of the above tumor strand. The pathological tissues are *S T*, sieve tubes; *C*, cambium; *Tr*, trachei; *Sp*, spiral vessels..... 182

PLATE XXII. Teratoid crowngalls produced in a tobacco plant by inoculating *Bacterium tumefaciens* into the leaf axils..... 182

PLATE XXIII. Teratoid crowngalls produced in tobacco leaves with the hop strain of *Bacterium tumefaciens* by local inoculations—that is, inoculation in places where shoot anlage are not known to exist. Fig. A.—Portion of an upper leaf showing four shoot-bearing tumors growing from upper surface of the inoculated midrib. Fig. B.—Same as *A*, but the leaf reversed and the midrib stripped of its blade to show two other shoot-bearing tumors which have developed from its under surface. Fig. C.—From middle of another leaf on the same plant as *A*, but further magnified and photo made on an orthochromatic plate to show the pale green character of the shoot as contrasted with the dark green of the surrounding leaf..... 182

HYPERASPIS BINOTATA, A PREDATORY ENEMY OF THE TERRAPIN SCALE

PLATE XXIV. *Hyperaspis binotata:* Fig. 1.—Male, showing the characteristic markings. Fig. 2.—Female, showing the dorsal view. Fig. 3.—Eggs and a second-instar larva. *a*, Second-instar larva as disclosed by displacing the host; *b*, larvæ of the terrapin scale, *Eulecanium nigrofasciatum; c*, a displaced scale; *d*, eggs "in situ"; *e*, egg. Fig. 4.—First-instar larva. Fig. 5.—Method of attacking the mature scales during the third and fourth instars.. 204

PLATE XXV. *Hyperaspis binotata:* Fig. 1.—Mature larva as it appears when attacking the leaf-attached larvæ of the terrapin scale, *Eulecanium nigrofasciatum*. Fig. 2.—Ventral view of mature larva. Fig. 3.—Dorsal view of pupa, showing the last larval molt skin and the depressed segmented area. Fig. 4.—Ventral view of pupa..... 204

TESTS OF THREE LARGE-SIZED REINFORCED-CONCRETE SLABS UNDER CONCENTRATED LOADING

PLATE XXVI. Fig. 1.—Load-applying and load-measuring apparatus for testing reinforced-concrete slabs, showing set-up for 4-point loading. Fig. 2.—Load-measuring apparatus and hydraulic jack for testing reinforced-concrete slabs..... 234

OCCURRENCE OF STERILE SPIKELETS IN WHEAT

PLATE XXVII.—Comparison of the number of sterile spikelets on bearded and beardless varieties of wheat..... 250

EGG AND MANNER OF OVIPOSITION OF LYCTUS PLANICOLLIS

PLATE XXVIII. *Lyctus planicollis:* Fig. 1.—Outline of the egg, showing strand attachment. Fig. 2.—Greatly enlarged view of end of egg, showing granular appearance. Fig. 3.—Larva within egg, ready to hatch. Fig. 4.—Sketch of egg in pore of wood on radial section of green-ash ladder-rung stock, showing longitudinal striæ..... 276

Page

PLATE XXIX. *Lyctus planicollis:* Larval burrows in an ash shovel handle.... 276
PLATE XXX. *Lyctus planicollis:* Pupal cells in "powder-posted" white-ash shovel handle..... 276
PLATE XXXI. *Lyctus planicollis:* Exit holes of adults in ash shovel handles.. 276

HYPODERMA DEFORMANS, AN UNDESCRIBED NEEDLE FUNGUS OF THE WESTERN YELLOW PINE

PLATE XXXII. Fig. 1.—Needles of *Pinus ponderosa* infected with *Hypoderma deformans*, showing the apothecia. Fig. 2.—Branches of *Pinus ponderosa* deformed and broomed by *Hypoderma deformans*. Fig. 3.—A branch of *Pinus ponderosa*, showing how it will hang vertically when supporting a large broom caused by *Hypoderma deformans*..... 288

ORNIX GEMINATELLA, THE UNSPOTTED TENTIFORM LEAF MINER OF APPLE

PLATE XXXIII. *Ornix geminatella* Pack.: Fig. 1.—Moth expanded. Fig. 2—Moth at rest on leaf. Fig. 3.—Egg on lower surface of leaf; also tunnel made by miner on leaving the egg. Fig. 4.—Dorsal view of first larval stage; below, side view of head and thorax. Fig. 5.—Dorsal view of second larval stage. Fig. 6.—Side view of second larval stage. Fig. 7.—Dorsal view of third larval stage, showing edge of thoracic legs. Fig. 8.—Dorsal view of fourth larval stage. Fig. 9.—Side view of fourth larval stage. Fig. 10.—Ventral view of pupa. Fig. 11.—Dorsal view of same. Fig. 12.—Lower surface of leaf with numerous partly developed mines; also two cocoons, one exposed. Fig. 13.—Portion of leaf showing a mine in process of development. Fig. 14.—A small twig showing leaves badly curled and injured by numerous mines. Fig. 15.—Leaf much distorted with 10 mines almost completed; also one cocoon appears at the tip of the leaf..... 296

A WESTERN FIELDROT OF THE IRISH POTATO TUBER CAUSED BY FUSARIUM RADICICOLA

PLATE XXXIV. Fig. 1, 2, 3, 4.—Types of stem-end blackrot, lenticel rot, and eye rot in Idaho Rural potato tubers. Fig. 5, 6.—Longitudinal and cross sections of an Idaho Rural tuber infected with blackrot..... 310
PLATE XXXV. Fig. 1.—Netted Gem potato tuber infected with jelly-end rot. Fig. 2.—Stem-end view of a Netted Gem tuber infected with jelly-end rot. Fig. 3.—Longitudinal section of a Netted Gem tuber infected with jelly-end rot. Fig. 4.—Idaho Rural tuber infected with stem-end and lenticel blackrot, after having been kept 10 days in a moist chamber. Fig. 5.—Idaho Rural tuber infected with lenticel blackrot after having been kept in a moist chamber for 10 days..... 310
PLATE XXXVI. Fig. 1, 2.—Stem-end blackrot produced by stem-end punctures with a needle carrying *Fusarium radicicola*. Netted Gem and Idaho Rural potato tubers. Fig. 3.—Lenticel blackrot produced by spraying the tuber with a spore suspension of *F. radicicola*. Netted Gem tuber. Fig. 4—Same tuber as shown in figure 3; after having been kept a few days longer in the moist chamber. Fig. 5.—Stem-end blackrot produced by spraying an Idaho Rural tuber with a spore suspension of *F. radicicola*. Fig. 6.—Stem-end blackrot produced by the inoculation of the tuber stolon. Idaho Rural tuber. Fig. 7.—Blackened vascular system produced by the inoculation of the tuber stolon. Idaho Rural tuber..... 310

Page

PLATE XXXVII. Fig. 1, 2, 3.—Stem-end and lenticel blackrot produced by spraying the growing tubers witn a spore suspension of *Fusarium radicicola*. Idaho Rural potato tubers. Fig. 4.—Eye infection produced by spraying the growing tuber with a spore suspension of *F. radicicola*. Netted Gem tuber. Fig. 5, 6, 7.—Stem-end blackrot produced by the inoculation of the stolons of growing Idaho Rural tubers. Fig. 8.—Stem-end rot of Netted Gem tuber produced by inoculating the stolon of the growing tuber..... 310

COMPARATIVE STUDY OF THE ROOT SYSTEMS AND LEAF AREAS OF CORN AND THE SORGHUMS

PLATE XXXVIII. Fig. 1.—Method used in isolating root systems in the field. View of two soil prisms ready for washing. Fig. 2.—Method used in isolating root systems. Fig. 3.—Method of washing used in the isolation of the root systems..... 332

PLATE XXXIX. Fig. 1.—Root system of a corn plant that had reached a height of 3 feet 6 inches. Fig. 2.—Root systems of two corn plants with a height of 1 foot 6 inches. Fig. 3.—Root system of a Dwarf milo plant at the age of 4 weeks. Fig. 4.—Root systems of two Blackhull kafir plants 1 foot in height..... 332

PLATE XL. Fig. 1.—Root systems of two mature corn plants. Fig. 2.—Root system of a corn plant at the time of "shooting"..... 332

PLATE XLI. Fig. 1.—Root systems of two Blackhull kafir plants at the time they had reached a height of 6 feet and were blooming. Fig. 2.—Root system of two Dwarf milo plants at the time the seed was in the milk stage. 332

PLATE XLII. Fig. 1.—Portion of a primary root of Pride of Saline corn, showing the number and relative size of the secondary roots. Fig. 2.—Portions of the primary roots of Blackhull kafir, showing the number and relative size of the secondary roots..... 332

PLATE XLIII. Fig. 1.—Pride of Saline corn, Dwarf milo, and Blackhull kafir plants, showing their relative leaf and sheath areas at 4 weeks of age. Fig. 2.—Pride of Saline corn, Dwarf milo, and Blackhull kafir plants, showing their relative leaf and sheath areas at 6 weeks of age..... 332

PLATE XLIV. Fig. 1.—Pride of Saline corn, Dwarf milo, and Blackhull kafir plants, showing their relative leaf and sheath areas at 8 weeks of age. Fig. 2.—Pride of Saline corn, Dwarf milo, and Blackhull kafir plants, showing their relative leaf and sheath areas at 10 weeks of age..... 332

SILVER-SCURF OF THE IRISH POTATO CAUSED BY SPONDYLOCLADIUM ATROVIRENS

PLATE XLV. Fig. 1.—Potato tubers showing shriveling and a silvery appearance caused by *Spondylocladium atrovirens*. Fig. 2.—Tuber naturally infected by *S. atrovirens*, showing the segregated area type of infection. Fig. 3.—Immature potato tuber artificially inoculated with conidia of *S. atrovirens*, July, 1913, at Houlton, Me..... 350

PLATE XLVI. Fig. 1.—Photomicrograph of *Spondylocladium atrovirens* on corn-meal agar, showing method of development of conidiophores and conidia in the early stages. Fig. 2.—Photomicrograph of *S. atrovirens* in hanging-drop culture, showing development of conidiophore and conidia in mature stages. Fig. 3.—Negative heliotropism of *S. atrovirens* on corn-meal agar exposed on one side to daylight from April 8 to April 24, 1915, in laboratory at room temperature..... 350

PLATE XLVII. Effect of temperature upon mycelial development of *Spondylocladium atrovirens* in pure culture on corn-meal agar at end of four weeks..... 350

Page

PLATE XLVIII. Contact infection. A part of the new tubers becoming infected with *Spondylocladium atrovirens* by means of contact with the infected mother tuber..... 350

PATHOLOGICAL HISTOLOGY OF STRAWBERRIES AFFECTED BY SPECIES OF BOTRYTIS AND RHIZOPUS

PLATE XLIX. *A–H*, Strawberry cells attacked by *Botrytis* sp.: *A*, Hypha growing partly between and partly within strawberry cells; *B*, hyphæ inside strawberry cells in which remnants of the protoplasm may still be distinguished; *C*, hypha passing from one cell into another across a short intercellular space; *D*, *E*, *F*, *G*, *H*, hyphæ entering cells in various ways (in *G* the hypha has pushed back a portion of the cell wall before breaking through). *I–M*, Strawberry cells attacked by *Rhizopus* sp.: *I*, Hypha growing between the epidermis and the adjacent layer of storage cells; *J*, hypha growing over the surface of the strawberry; *K*, hyphæ growing underneath the epidermal layer and between the storage cells; *L*, *Rhizopus* sp. growing between epidermal cells; *M*, germinating spore in cavity formed by a seed..... 366

PLATE L. Strawberry cells attacked by *Rhizopus* sp. *A*, Normal storage cell of strawberry; *B*, storage cell showing a slight contraction of the protoplasm; *C*, *D*, *E*, *F*, *G*, progressive contraction of protoplasm of host cells near hyphæ; *H*, *I*, *J*, strawberry cells near hyphæ in which the cell wall has crumpled with the contraction of the protoplasm; *K*, *M*, hyphæ inside cells; *L*, hyphæ growing between cells of the strawberry; *K*, *L*, *M* are drawn from berries which had been rotted in the desiccator..... 366

LIFE HISTORIES AND METHODS OF REARING HESSIAN-FLY PARASITES

PLATE LI. Fig. 1.—Egg of *Eupelmus allynii*. Fig. 2.—Egg of *Eupelmus allynii* in situ. Figs. 3, 4.—Pupa of *Eupelmus allynii*. Fig. 5.—Egg of *Merisus destructor*. Fig. 6.—Pupa of *Merisus destructor*. Fig. 7.—Egg of *Micromelus subapterus*. Fig. 8.—Pupa of *Micromelus subapterus*..... 382

PLATE LII. Fig. 1.—Mandibles of full-grown larva of *Eupelmus allynii*. Fig. 2.—Larva of *Eupelmus allynii*. Fig. 3.—Mandibles of full-grown larva of *Merisus destructor*. Fig. 4.—Larva of *Merisus destructor*. Fig. 5.—Mandibles of full-grown larva of *Micromelus subapterus*. Fig. 6.—Larva of *Micromelus subapterus*..... 382

TRANSMISSION AND CONTROL OF BACTERIAL WILT OF CUCURBITS

PLATE LIII.—Two wilted cucumber plants which contracted bacterial wilt at beetle gnawings of the leaves marked *x*. Three healthy, uninjured plants in same hill are also shown..... 434

PLATE LIV.—Plots in field 1, East Marion, Long Island, N. Y., 1915. Fig. 1.—Plot sprayed with Bordeaux mixture and lead arsenate, beginning June 25. Fig. 2.—Plot sprayed with Bordeaux mixture and lead arsenate, beginning July 19, after most of the striped-beetle injury had occurred. Fig. 3.—Plot sprayed with Bordeaux mixture and lead arsenate, beginning July 27. Fig. 4.—General view of field, showing cages and meteorological-instrument shelter..... 434

CORRELATED CHARACTERS IN MAIZE BREEDING

PLATE LV.—Typical plant of the Waxy Chinese variety of maize, showing numerous tassel branches, erect leaf blades, one-sidedness, and curved tassel..... 454

Page

PLATE LVI. Fig. 1.—Uppermost leaf sheaths of Chinese maize plant, showing the one-sided arrangement of leaf blades and absence of hairs. Fig. 2.—Leaf sheath of the Waxy Chinese variety of maize, showing the transverse lines and absence of hairs 454

PLATE LVII.—A plant of the Esperanza variety of maize, showing the drooping leaves, few tassel branches, and elongated internodes characteristic of the variety 454

PLATE LVIII.—Leaf sheaths of the Esperanza variety of maize, showing the maximum development of tuberculate hairs 454

PLATE LIX.—A leaf sheath of a second-generation hybrid maize plant 454

PLATE LX.—A first-generation plant of Chinese×Esperanza maize hybrid 454

PLATE LXI.—A second-generation plant of a Chinese×Esperanza maize hybrid 454

PLATE LXII.—A second-generation plant of a Chinese×Esperanza maize hybrid 454

PLATE LXIII.—A second-generation plant of a maize hybrid, showing the "goose-neck" character that appeared for the first time in this hybrid 454

ALEYRODIDAE, OR WHITE FLIES ATTACKING THE ORANGE, WITH DESCRIPTIONS OF THREE NEW SPECIES OF ECONOMIC IMPORTANCE

PLATE LXIV.—*Aleurocanthus woglumi:* Eggs, larvæ, and pupa cases on orange leaves 472

PLATE LXV.—*Aleurocanthus woglumi:* Fig. 1.—Colony on an orange leaf. Fig. 2.—Eggs and pupa cases 472

PLATE LXVI.—Fig. 1.—*Dialeurodes citri:* Pupæ. Fig. 2.—Male and female adults of an aleyrodid. Fig. 3.—*Aleurolobus marlatti* 472

PLATE LXVII. *Aleurothrixus howardi:* Larvæ and pupa cases on an orange leaf 472

PLATE LXVIII. *Aleurothrixus porteri:* Larvæ and pupa cases on *Myrtus* sp. 472

PLATE LXIX. Fig. 1.—*Tetraleurodes mori*, var. *arizonensis:* Larvæ and pupa cases on an orange leaf. Fig. 2.—*Tetraleurodes mori:* Pupa cases on a mulberry leaf 472

RELATIVE WATER REQUIREMENT OF CORN AND THE SORGHUMS

PLATE LXX. Fig. 1.—General view of the screened inclosure and the scale house. Fig. 2.—Method of moving the cans. Fig. 3.—General view of the plant shelter and the surrounding country at Garden City, Kans 484

PLATE LXXI. Fig. 1.—Dwarf milo, grown May 22 to September 3, 1915. Fig. 2.—Dwarf Blackhull kafir, grown May 22 to September 11, 1915. Fig. 3.—Feterita, grown May 22 to September 6, 1915 484

PLATE LXXII. Fig. 1.—Sudan grass, grown May 22 to September 14, 1915. Fig. 2.—Pride of Saline corn, grown May 22 to August 25, 1915. Fig. 3.—Blackhull kafir, grown May 22 to September 18, 1915. Fig. 4.—Method of sealing the lids with tape and the wax seal around the plants 484

AVAILABILITY OF MINERAL PHOSPHATES FOR PLANT NUTRITION

PLATE LXXIII. Effect of varying quantities of Tennessee brown rock phosphates on plant growth: Fig. 1.—Spring wheat. Fig. 2.—Sixty-Day oats .. 514

PLATE LXXIV. Effect of varying quantities of Tennessee brown rock phosphate on plant growth: Fig. 1.—Barley. Fig. 2.—Timothy 514

PLATE LXXV. Effect of varying quantities of Tennessee brown rock phosphate on plant growth: Fig. 1.—Cowpeas. Fig. 2.—Soybeans. Fig. 3.—Red clover. Fig. 4.—Alfalfa 514

PLATE LXXVI. Effect of different kinds of mineral phosphate applied in different quantities for red clover 514

Page

PLATE LXXVII. Cowpeas, showing the comparative effect of Tennessee brown rock phosphate alone and in combination with dextrose..... 514

PLATE LXXVIII. Cowpeas, showing the comparison of their growth when treated with Tennessee brown rock phosphate, phosphate and dextrose, and phosphate, dextrose, and calcium carbonate..... 514

PLATE LXXIX. Effect of different substances on the growth of cowpeas: Fig. 1.—Growth after the addition of varying quantities of raw rock. Fig. 2.—Growth after the addition of dextrose and soluble phosphate..... 514

PLATE LXXX. Effect of various substances and combinations on the growth of cowpeas: Fig. 1.—Effect of adding lime, phosphate rock, dextrose and lime, and phosphate rock, dextrose, and lime to the soil. Fig. 2.—Effect of adding nothing, lime, phosphate rock, and phosphate rock and lime to the soil..... 514

STORAGE-ROTS OF ECONOMIC AROIDS.

PLATE LXXXI. Fig. 1.—A dasheen corm (*Colocasia esculenta*) showing Java blackrot produced by *Diplodia tubericola*. Fig. 2.—A corm of *Alocasia sp.* showing Java blackrot produced by *D. tubericola*. Fig. 3.—A dasheen tuber partially decayed by *Sclerotium rolfsii*..... 572

PLATE LXXXII. Fig. 1.—A tuber of *Colocasia esculenta* showing a powdery grayrot caused by *Fusarium solani*. Fig. 2.—A tuber of *Xanthosoma sagittifolium* showing partial decay by *Fusarium solani*. Fig. 3.—A tuber of *C. esculenta* softened throughout by *Sclerotium rolfsii*..... 572

PLATE LXXXIII. A corm of *Colocasia esculenta* from Brooksville, Fla., mostly rotted away by *Bacillus carotovorus*..... 572

INFLUENCE OF CALCIUM AND MAGNESIUM COMPOUNDS ON PLANT GROWTH

PLATE LXXXIV. Fig. 1.—Growth of wheat in sand containing varying quantities of calcium and magnesium. Fig. 2.—Growth of alfalfa in sand containing varying amounts of calcium and magnesium..... 620

PLATE LXXXV. Fig. 1.—Growth of soybeans following a crop which had already absorbed most of the readily available calcium and magnesium. Fig. 2.—Growth of soybeans in soil treated with magnesium..... 620

PLATE LXXXVI. Fig. 1.—Comparative growth of soybeans in brown silt loam and dolomite, showing that the loam would have been improved by the addition of some limestone or dolomite. Fig. 2.—Soybeans in sand treated with magnesium, showing that their growth increases inversely with the quantity of magnesium applied as sulphate..... 620

PLATE LXXXVII. Comparative root production of wheat grown in the chlorids, sulphates, and carbonates of magnesium and calcium..... 620

PLATE LXXXVIII. Fig. 1.—Comparative growth of wheat in sand and in dolomite. Fig. 2.—Comparative growth of wheat in magnesium chlorid and magnesium sulphate..... 620

LARVAL CHARACTERS AND DISTRIBUTION OF TWO SPECIES OF DIATRAEA

PLATE LXXXIX. Fig. 1.—*Diatraea saccharalis crambidoides:* Larva, summer form, dorsal view. Fig. 2.—*D. zeacolella:* Larva, summer form, dorsal view. Fig. 3.—*D. saccharalis crambidoides:* Larva, summer form, side view. Fig. 4.—*D. seacolella:* Larva, summer form, side view. Fig. 5.—*D. saccharalis crambidoides:* Larva, winter form, dorsal view. Fig. 6.—*D. zeacolella:* Larva, winter form, dorsal view..... 626

THE DISEASE OF POTATOES KNOWN AS "LEAK"

Page

PLATE XC. Potatoes affected with potato leak: Fig. 1, 2.—Natural infection from fork wound. Fig. 3.—Rot produced by inoculation with *Pythium debaryanum*. Fig. 4.—Rot produced by inoculation with *Rhizopus nigricans*..... 640

SOME PROPERTIES OF THE VIRUS OF THE MOSAIC DISEASE OF TOBACCO

PLATE XCI. Livingstone atmometer porus cup as used for filtration..... 674

LIFE CYCLES OF THE BACTERIA

PLATE A. Fig. 1.—Azotobacter 11. Mannite-nitrate solution, 5 days old. Types A and Lα. Some cells in conjunction. Fig. 2.—Azotobacter 21. Contact preparate from a colony on mannite agar, 4 days old. Types A, L. Most cells in conjunction. Fig. 3.—Azotobacter 23. Contact preparate from a colony on mannite-agar, 4 days old. Types A, B, I, Kα, and many conjunct cells. Fig. 4.—Azotobacter 13. Jannite-nitrate solution, 17 days old. Type Kλ. Fig. 5.—Azotobacter 14. Mannite-nitrate solution, 5 days old. Type B forming I. Fig. 6.—Azotobacter 8. Beef bouillon. Type B forming types I and J..... 702

PLATE B. Fig. 7.—Azotobacter 21. Mannite-agar colony, 4 days old. Type C forming types D and I. Fig. 8.—Azotobacter 22. Mannite-agar colony, 4 days old. Type C forming D, also A in conjunction. Fig. 9.—Azotobacter 11. From a filter paper strip in mannite-peptone solution, 16 days old. Types A and B forming D. Fig. 10.—Azotobacter 3. Mannite-peptone solution, 24 days old. Types L and M forming D. Fig. 11.—Azotobacter 11. Mannite-peptone solution, 16 days old. Type D (stained) resulting from Type C. Fig. 12.—Azotobacter 6. From condensation water of mannite-agar slant, 1 day old. Type D (unstained) containing regenerative units..... 702

PLATE C. Fig. 13.—Azotobacter 24. Mannite-nitrate solution kept 5 days after having been heated 1 minute at 96° C. Types I and F developing from D. Some I germinating in conjunct stage and inclining to form spores. Fig. 14.—Azotobacter 1. Mannite-nitrate solution, 10 days old. Types B, Kβ, E, and Fα developing from stained and unstained type D. Fig. 15.—Azotobacter 15. From condensation water of a mannite-nitrate agar slant, 2 days old. Types Fα and Fβ developing from type D. Fig. 16.—Azotobacter 17. Mannite-soil-extract agar, 2 months old. Type E, Fα, Kφ, and G developing from type D. Fig. 17.—Azotobacter 17. Mannite-nitrate agar, 10 days old. Preparate treated with hot aqueous fuchsin. Type G, partially dissolved; also Kφ. Fig. 18.—Azotobacter 7. Mannite-soil-extract solution, 14 days old. Type H forming D..... 702

PLATE D. Fig. 19.—Azotobacter 2. Mannite-nitrate agar, 23 days old. Spores forming type D. Fig. 20.—Azotobacter 2. Mannite nitrate-agar, 6 days old. Types L and F, endospores and exospores and dissolving of spores to type D. Fig. 21.—Azotobacter 18. From a filter paper strip in mannite solution, 25, days old. Type L with gonidia, forming B (type Jλ). Fig. 22.—Azotobacter 7. Mannite-soil-extract agar, 2 months old. Types E and F forming B. Fig. 23.—Azotobacter 7. Mannite-soil-extract agar, 2 months old. Type B, formed by types E and F, germinating to type G. Fig. 24.—Azotobacter 7. Mannite-soil-extract agar, 2 months old. Type Kγ..... 702

55854°—16——2

Page

PLATE E. Fig. 25.—*Bacillus subtilis* (No. 31). Beef agar, 2 days old. Types I and D formed by spores. Fig. 26.—*Bacillus subtilis* (No. 31). Beef agar, 6 days old. Formation of type I. Fig. 27.—*Bacillus subtilis* (No. 31). Beef agar, 8 days old. Type I forming H and stained D. Spores forming unstained type D. Fig. 28.—Yellow bacillus (No. 41). Peptone-glycerin solution, 2 days old. Type I germinating from D, stretching to type L. Fig. 29.—*Bacterium bulgaricum* (No. 49). Whey-yeast agar, 6 days old at 40° C. Types C, D, E, F, G, I, and K. Fig. 30.—*Bacterium fluorescens* (No. 40). Ammonium-citrate-glycerin solution, 11 days old. Types D and H. 702

PLATE F. Fig. 31.—*Sarcina flava* (No. 43). Beef agar, 1 day old. Type I in conjunction and forming D. Fig. 32.—*Streptococcus lactis* (No. 48). Peptone lactose solution, 5 days old. Type D, with regenerative units, forming type I. Fig. 33.—*Streptococcus lactis* (No. 48). Milk, 3 days old. Types D and I in casein. Fig. 34.—*Bacillus radicicola* (No. 39). Types D and I. Preparate made from a root nodule in 1908. Fig. 35.—*Spirillum* sp. from Great Salt Lake (No. 46). Beef broth plus 3 per cent of sodium chlorid, 14 days old. Budding and branching forms; stained and unstained regenerative bodies. Some cells in conjunction. Fig. 36.—*Spirillum* sp. from Great Salt Lake (No. 46). Beef broth plus 3 per cent of sodium chlorid, 14 days old. Type I germinating 702

PLATE G. Fig. 37.—*Micrococcus candicans* from soil (No. 45). Ammonium-citrate-glycerin solution, 6 days old. Irregular, thick-walled type I. Fig. 38.—*Micrococcus candicans* from milk (No. 44). Ammonium-citrate-glycerin solution, 2 days old. Irregular, thick-walled type I. Fig. 39.—Yellow bacillus (No. 41). Beef agar, 1 day old. Budding gonidia, formation and germination of type I. Fig. 40.—*Bacterium fluorescens* (No. 40). Ammonium-citrate-glycerin solution, 2 days old. Budding gonidia, formation and germination of type I. Fig. 41.—*Bacterium fluorescens* (No. 40). Beef agar, 4 days old. Filterable gonidia germinating. Fig. 42.—*Bacterium fluorescens* (No. 40). Beef agar, 4 days old. Types D and F formed by filterable gonidia. Dark field 702

A RESPIRATION CALORIMETER, PARTLY AUTOMATIC, FOR THE STUDY OF METABOLIC ACTIVITY OF SMALL MAGNITUDE

PLATE XCII.—General view of the respiration calorimeter: *A*, Chamber inclosed in heat-insulating cover. *B*, Tension equalizer to maintain atmospheric pressure in the air of the chamber. *C*, Absorber table. *D*, Rotary pump to maintain air circulation. *E*, Motor to drive pump. *F*, Bottles containing sulphuric acid to remove water vapor from circulating air. *G*, Large U-tube, containing soda-lime to remove carbon dioxid from the air. *H*, Bottle containing sulphuric acid to catch the water vapor from the soda-lime. *I*, Bottle containing cotton to catch sulphuric acid vapor. *J*, Small absorbers for determining water vapor and carbon dioxid in residual air. *K*, Meter to measure the sample of residual air. *L*, Reservoir to maintain a constant pressure of water in the heat absorber in the chamber. *M*, Tank to catch water flowing from the heat absorber. *N*, Pump to raise water from the tank to the reservoir. *O*, Devices for automatically controlling and recording temperatures 720

Page

PLATE XCIII.—Chamber with part of outer covering removed: *A*, Double metal wall chamber. *B*, Heat-insulating outer cover. *C*, Window to chamber. *D*, Outlet providing passage for pipes, wires, etc., through the walls of the chamber. The exterior ends of the resistance thermometers for ingoing and outgoing water are seen projecting from the outlet. *E*, Removable top of chamber. *F*, Device for heating the air entering the respiration chamber. *G*, Small pipe carrying water for cooling the outer metal wall of the chamber. *H*, Electric-resistance wire carrying current for heating the outer wall..... 720

PLATE XCIV.—Apparatus connected with the respiration calorimeter: *A*, Tension equalizer. *B*, Mixing bottle for equalizing the temperature of water entering the heat absorber. *C*, Device for heating air entering the respiration chamber. *D*, Preheater, and *E*, final heater, for raising the temperature of water entering the heat absorbers. There is an electric-heating coil in the lower half and an electric-resistance thermometer in the upper half of the final heater. *F*, Temperature indicator comprising part of the apparatus for controlling the temperature of the water entering the heat absorber. *G*, Multiple-point switch for connecting the resistance thermometers for the metal walls and air of the chamber with the Wheatstone bridge for measuring their temperatures. *H*, Tube conducting air from the respiration chamber to the rotary air pump. *I*, Tube conducting air from the purifying devices to the respiration chamber..... 720

PLATE XCV.—Devices for controlling and recording temperatures: *A*, Mechanism for shifting the contact on the rheostats controlling the current for heating the outer walls of the calorimeter chamber and the ingoing air. *B*, Ratio coils for the four bridges governing the action of the shifting mechanism *A* are combined in this box, together with means for checking the constancy of the resistance of the coils and for correcting slight inequalities in them and also to compensate for small differences in the pair of resistance thermometers forming the other arms of each bridge. *C*, Mechanism for shifting the contact on the rheostat controlling the current in the heating coil in the final heater, shown at *E* in Plate XCIV. *D*, Temperature-difference recorder (self-balancing Wheatstone bridge) for continuously recording the difference between the temperature of the water entering and that leaving the heat absorber. *E*, "Check box" containing the ratio coils of the bridge for temperature difference measurements and coils for extending the range of differences measured, with means for checking the constancy of the resistances of the coils and the accuracy of the recorder readings and also for compensating for slight differences in the resistance of the thermometer coils when they are at the same temperature..... 720

MOTTLE-LEAF OF CITRUS TREES IN RELATION TO SOIL CONDITIONS

PLATE H.—Various stages in mottle-leaf of the orange..... 740

PLATE XCVI.—Orange leaves showing mottle-leaf..... 740

PLATE XCVII.—A more advanced stage of mottle-leaf of orange, showing the reduction in the size of the leaves..... 740

VEGETATIVE SUCCESSION UNDER IRRIGATION

PLATE XCVIII.—Rock Creek Valley, near Rock River Station..... 760

PLATE XCIX.—A nearer view of the bench slope; the same tree shown in Plate C, figure 1..... 760

PLATE C. Fig. 1.—Where upland and lowland meet. Fig. 2.—Characteristic draw; the stream valley beyond. Lupin, wheat-grass, white sage, and gaillardia in the foreground..... 760

Page

PLATE CI. Fig. 1.—The bench. The course of Rock Creek is indicated by the distant trees. Fig. 2.—Part of a reservoir on the Rock Creek ranch..... 760

PLATE CII. Fig. 1.—Lupin recessive and cat's-foot becoming dominant. Gay's sedge subphase in background. Fig. 2.—Wheat-grass phase..... 760

PLATE CIII. Fig. 1.—Squirrel-tail phase. A few grindelias in the foreground. Fig. 2.—Rush-sedge phase (the darker areas) replacing wheat-grass phase. 760

PLATE CIV. Fig 1.—Hair-grass phase. Fig. 2.—Natural meadow..... 760

PLATE CV. Fig. 1.—Field of oats on bench. Cat's-foot and other upland plants in foreground. Fig. 2.—Alfalfa field one year after sowing. Cat's-foot and bench grasses in foreground..... 760

AGRICULTURAL VALUE OF IMPERMEABLE SEEDS

PLATE CVI. Fig. 1.—A row of alsike clover from impermeable seeds between two rows from permeable seeds. Fig. 2.—A row of white clover from impermeable seeds between two rows from permeable seeds..... 796

LIFE-HISTORY STUDIES OF CIRPHIS UNIPUNCTA, THE TRUE ARMY WORM

PLATE CVII.—*A*, Cages for rearing *Cirphis unipuncta; B*, leaves glued together after the eggs have been deposited; *C*, characteristic leaves partly eaten by first-instar larvæ; *D*, full-grown larva; *E* and *F*, characteristic pupal cells. 812

CONTROL OF THE POWDERY DRYROT OF WESTERN POTATOES CAUSED BY FUSARIUM TRICHOTHECIOIDES

PLATE CVIII. Fig. 1.—A potato tuber infected with powdery dryrot, showing the wrinkled condition of skin due to the decay of underlying tissues. Fig. 2.—A potato tuber infected with powdery dryrot. Advanced stage. Fig. 3.—Section through a potato tuber infected with powdery dryrot, showing the internal cavities filled with the mycelium and the spores of the fungus..... 832

THERSILOCHUS CONOTRACHELI, A PARASITE OF THE PLUM CURCULIO

PLATE CIX. *Thersilochus conotracheli:* Fig. 1.—Adult female. Fig. 2.—*a*, Adult male; *b*, side view of abdomen..... 856

APHIDOLETES MERIDIONALIS, AN IMPORTANT DIPTEROUS ENEMY OF APHIDS

PLATE CX. *Aphidoletes meridionalis:* Fig. 1.—Adult female: *a*, Antenna of male, showing structure; *b*, tip of male abdomen. Fig. 2.—Larva attacking a pea aphis..... 888

PROGRESSIVE OXIDATION OF COLD-STORAGE BUTTER

PLATE CXI.—Gas apparatus used in the extraction and analysis of the air confined in butter..... 952

STUDIES ON THE PHYSIOLOGY OF REPRODUCTION IN THE DOMESTIC FOWL.—XV. DWARF EGGS

PLATE CXII.—A collection of dwarf eggs with a normal egg in the center of the group..... 1042

PLATE CXIII.—Fig. 1.—Ovarian follicles (*b–f*) and the dwarf egg *a* from case 27. Fig. 2.—Shell of a compound egg which was composed of two albumen masses partly separated at the level of the seam in the shell by an incomplete egg membrane. Fig. 3.—Dwarf egg containing a mass of yolk not in a yolk membrane, but separated from the albumen by a membrane-like layer of chalazal threads. Fig. 4.—Dwarf egg formed around an artificial yolk of agar which was inserted into the oviduct. *a*, Complete egg; *b*, agar yolk..... 1042

α-CROTONIC ACID, A SOIL CONSTITUENT

Page

PLATE CXIV. Fig. 1.—α-Crotonic acid from soil. Fig. 2.—Synthetic α-crotonic acid 1046

TEXT FIGURES

CLIMATIC CONDITIONS AS RELATED TO CERCOSPORA BETICOLA

FIG. 1. *Cercospora beticola: A*, Section of overwintered sugar-beet leaf showing embedded sclerotia-like body, *a*, with a mass of old conidiophores, *b*, from which a new conidium, *c*, was produced. *B*, Production of rather typical conidiophores, *b*, and conidia, *c*, from a sclerotia-like mass, *a*, taken from overwintered host material and placed in hanging-drop cultures 22

2. Curves of the maximum and minimum soil and air temperatures for the period from December 5, 1912, to March 13, 1913, at Rocky Ford, Colo., and air temperatures from December 5, 1913, to March 13, 1914, at Madison, Wis., together with the periods that snow covered the ground 28

3. Curves of the maximum and minimum soil and air temperatures for Rocky Ford, Colo., from March 13 to June 17, 1913, and for Madison, Wis., from March 13 to June 17, 1914 29

4. Curves of the maximum and minimum temperatures and relative humidities and the number of hours that the humidity remained above 60 from noon of the preceding to noon of the given day among sugar-beet plants and in the air 5 feet above the field, together with the field rainfall and irrigation records. June 11 to August 2, 1913, at Rocky Ford, Colo 32

5. Curves of the maximum and minimum temperatures among sugar-beet plants and at the Weather Bureau station, and the seasonal rainfall records at Madison, Wis., in 1914, and the number of hours that the humidity remained above 60 among the sugar-beet plants in the field at Madison, Wis., in 1914, and at Rocky Ford, Colo., in 1913 34

6. Curves of the leaf spot history series, showing the production of conidia on different dates from June 24 to September 19, 1913, at Rocky Ford, Colo 48

7. Curves of the maximum and minimum temperatures and humidities, the number of hours that the humidity remained above 60 from noon of the preceding to noon of the given day among the plants, and rainfall and irrigation records, taken in a medium-early sugar-beet field from June 10 to September 22, 1913, at Rocky Ford, Colo .. 51

8. Curves of the comparative production of conidia on the upper and lower surfaces of the leaf spots, representing series E, K, N, and G of Table V and figure 6. Rocky Ford, Colo., 1913 52

9. Curves of the 2-day average increases in the number of leaf spots per plant in a medium-early and an early sugar-beet field, from June 18 to September 19, 1913, at Rocky Ford, Colo 54

10. Curves of the maximum and minimum temperatures and relative humidities and the number of hours that the humidity remained above 60 from noon of the preceding to noon of the given day among the sugar-beet plants of a medium-early and an early field. August 2 to September 21, 1913, at Rocky Ford, Colo 58

CITRUS CANKER

Page

FIG. 1. Diagrammatic representation of young open type of Citrus canker of half the diameter of the one shown in figure 2. *pp*, Palisade parenchyma; *ue*, upper epidermis; *le*, lower epidermis; *d*, diseased tissues; *a*, air space arising from tensions due to the enlargement of cells and disintegration of tissues 72

2. Diagrammatic representation of canker on old Citrus leaf; *pp*, Palisade parenchyma; *ue*, upper epidermis; *le*, lower epidermis; *p*, pycnidium of *Phoma socia; d*, diseased tissues; *a*, air space arising from tensions due to the enlargement of cells and disintegration of tissues 73

Pseudomonas citri: a, Stained with carbol fuchsin; *b*, stained with Williams's flagellar stain (adapted from Hasse); *c*, stained with anilin gentian violet 76

4. Early stage of Citrus canker in cross section on a young leaf of seedling grapefruit 80

5. *Pseudomonas citri:* (*a*). In the mesophyll tissue and (*b*) in the palisade parenchyma 81

6. Drawing of a stained section of a natural canker on grapefruit 82

7. Cross section in outline of a spongy canker on the rind of a fruit of *Citrus decumana*, showing ruptured epidermis and hypertrophy of the rind tissues, the cells of which are loosely attached 83

8. *a*, Cross section of a pycnidium of *Phoma socia* from a grapefruit leaf. *b*, Germination of conidia of *Phoma socia* after 24 hours in water. *c*, Mycelium of this fungus in old cultures 86

DETERMINATION OF STEARIC ACID IN BUTTER FAT

FIG. 1. Exterior of constant-temperature crystallization tank 104

2. Interior of constant-temperature crystallization tank 105

LIFE HISTORY AND HABITS OF TWO NEW NEMATODES PARASITIC ON INSECTS

FIG. 1. *Diplogaster labiata: A*, Mating; *B*, mature female reared in water culture, *a*, lip region, *b*, esophagus, *c*, median bulb, *d*, cardiac bulb, *e*, intestine, *f*, ovaries, *g*, egg, *h*, genital pore, *i*, rectum, *k*, anus; *C*, mature male reared in water culture, *a*, lip region, *b*, esophagus, *c*, median bulb, *d*, cardiac bulb, *e*, intestine, *k*, anus, *m*, spicula; *D*, at time of hatching; *E*, female during process of molting; *F*, dead female with young nematode which hatched within her body 116

2. *A–H, Diplogaster labiata:* Development of the egg; *I*, Diplogaster aerivora: mature male reared in moist soil; *J, Diplogaster aerivora:* mature male reared in water culture, *a*, lip region, *b*, esophagus, *c*, median bulb, *d*, cardiac bulb, *e*, intestine, *k*, anus, *m*, spicula; *K, Diplogaster aerivora:* dead female with young which hatched within her body; *L, Diplogaster aerivora:* mating 117

3. *Diplogaster aerivora: A*, Form found in termite; *B*, at time of hatching; *C*, female reared in water culture, not quite mature, *a*, lip region, *b*, esophagus, *c*, median bulb, *d*, cardiac bulb, *e*, intestine, *f*, ovaries, *h*, genital pore, *i*, rectum, *k*, anus; *D*, mature female reared in moist soil; *E*, mature female reared in water culture; *F–M*, development of the egg 122

EFFECT OF PASTEURIZATION ON MOLD SPORES

FIG. 1. Curve of the number of species of molds surviving pasteurization of milk for 30 minutes at a series of temperatures 157

2. Curve of the number of species of molds surviving flash pasteurization at a series of temperatures 160

Page

FIG. 3. Curve of the number of species of molds surviving dry heat for 30 minutes at a series of temperatures. 164

EFFECT OF WATER IN THE RATION ON THE COMPOSITION OF MILK

FIG. 1. Grouping of cow and kind of ration fed cows 23, 24, 25, and 27. 173

HYPERASPIS BINOTATA, A PREDATORY ENEMY OF THE TERRAPIN SCALE

FIG. 1. Map showing the distribution in the United States of *Hyperaspis binotata*. 198

TESTS OF THREE LARGE-SIZED REINFORCED-CONCRETE SLABS UNDER CONCENTRATED LOADING

FIG. 1. Diagram illustrating the method of obtaining "effective width" in reinforced-concrete slab tests. 207
2. Diagram showing location of strain-gauge points on top of slab 835. 211
3. Diagram showing location of strain-gauge points on top of slab 930. 212
4. Diagram showing location of strain-gauge points on top and bottom of slab 934. 213
5. Concrete deformation curves for concentrated center load on slab 835. 214
6. Concrete deformation curves for slab 930. 215
7. Deformation curves for slab 934. 216
8. Deformation curves for slab 934, computed from first zero reading. 216
9. Concrete deformation curves for slab 835 with 2-point loading. 217
10. Concrete deformation curves for slab 934 with 2-point loading. 217
11. Steel deformation curves for slab 934 with 2-point loading. 218
12. Concrete deformation curves for slab 835 with 4-point loading. 219
13. Concrete deformation curves for slab 934 with 4-point loading. 219
14. Deflection curves for slab 934 on first application of load. 220
15. Deflection curves for slab 934 on second application of load. 221
16. Deflection curves for slab 934 with 2-point loading. 222
17. Diagram showing effect of breaking load on slab 835. 223
18. Diagram showing effect of breaking load on slab 930. 223
19. Diagram showing effect of breaking load on slab 934. 224
20. Concrete deformation curves for a 10,000-pound concentrated center load on slab 934. Variation in deformations plotted parallel to supports. 225
21. Concrete deformation curves for a 10,000-pound concentrated center load on slab 934. Variation in deformations plotted perpendicular to supports. 226
22. Concrete deformation curves for slab 934. Lateral deformations plotted both parallel and perpendicular to supports. 227
23. Iso-deformation lines for slab 934 under concentrated center load. 228
24. Concrete deformation curves for slab 934 under 40,000-pound 4-point loading. Deformations measured perpendicular to supports. Variation of deformations plotted parallel to the supports. 229
25. Concrete deformation curves for slab 934 under 40,000-pound 4-point loading. Deformations measured perpendicular to supports. Variation of deformations plotted perpendicular to supports. 230
26. Iso-deformation lines for slab 934 under 40,000-pound 4-point loading. Deformations measured perpendicular to supports. 231
27. Curves showing effective width versus load (concentrated center load). 231
28. Curve showing effective width versus thickness. 232

EFFECT OF CLIMATIC FACTORS ON THE HYDROCYANIC-ACID CONTENT OF SORGHUM

Page

FIG. 1. Curves showing the hydrocyanic-acid content of sorghum on the various plots 264
2. Curves showing the rate of growth of the sorghum on the various plots.. 265
3. Curves showing the precipitation, temperature, and humidity relations at the various experiment stations during the growing season of 1915.. 266
4. Curves showing the contemporary climatic conditions at the various plots, together with crop data and hydrocyanic-acid content 267

HYPODERMA DEFORMANS, AN UNDESCRIBED NEEDLE FUNGUS OF THE WESTERN YELLOW PINE

FIG. 1. A side view of two apothecia of *Hypoderma deformans* on needles of *Pinus ponderosa*, showing the longitudinal medial split 278
2. Asci, spores, and paraphyses of *Hypoderma deformans* 279
3. Cross section of an apothecium of *Hypoderma deformans* on a needle of *Pinus ponderosa*, showing mature asci with spores, the point of first rupture, and the tissues of the leaf most seriously affected by the mycelium of the fungus 280
4. The upper portion of a young ascus of *Hypoderma deformans*, showing the formation of the pore at the tip through which the spores are expelled 281

COMPARATIVE STUDY OF THE ROOT SYSTEMS AND LEAF AREAS OF CORN AND THE SORGHUMS

FIG. 1. Evaporation from a free water surface (tank) at Garden City, Kans., during the growing seasons of 1914 and 1915 314
2. Comparison of the leaf areas of Pride of Saline corn, Black-hull kafir, and Dwarf milo at four stages of the growth of these plants during the season of 1914 328
3. A graphic illustration of the sheath areas of Pride of Saline corn, Black-hull kafir, and Dwarf milo at four stages of the growth of these plants during the season of 1914 329

WOOLLY PEAR APHIS

FIG. 1. Comparative structure of antennæ and wax pores of *Eriosoma* spp.: *A*, distal segments of antenna of winged viviparous female of *E. pyricola*; *B*, distal segments of antenna of winged viviparous female of *E. ulmi*; *C*, distal segments of antenna of wingless viviparous female of *E. americanum*; *D*, distal segments of antenna of wingless viviparous female of *E. lanigerum*; *E*, distal segments of antenna of wingless viviparous female of *E. pyricola*; *F*, distal segments of antenna of winged viviparous female of *E. americanum*; *G*, distal segments of antenna of winged viviparous female of *E. lanigerum*; *H*, compound wax pore of *E. lanigerum*; *I*, compound wax pore of *E. pyricola*; *J*, distal segments of antenna of first instar wingless viviparous female of *E. pyricola* 359

STIMULATING INFLUENCE OF ARSENIC UPON THE NITROGEN-FIXING ORGANISMS OF THE SOIL

FIG. 1. Graph showing the action of five compounds of arsenic on nitrogen fixation in dry soil 392
2. Graph showing the effect of aeration on the nitrogen-fixing activity of soil containing compounds of arsenic 408

Page

FIG. 3. Graph showing the effect of heat on the nitrogen-fixing power of soil treated and not treated with arsenic 409
4. Effect of various arsenic compounds in the ratio of 400 parts of the compound to 1,000,000 parts of soil on the activity of various soil organisms 411
5. Graph showing parts per million of various arsenic compounds in the soil at which the greatest stimulation occurred 412

TRANSMISSION AND CONTROL OF BACTERIAL WILT OF CUCURBITS

FIG. 1. Comparison of the amount of wilt with striped-beetle prevalence and with meteorological phenomena in three fields, East Marion, Long Island, N. Y., season of 1915 421
2. Comparison of relative wilt control of Bordeaux mixture plus lead arsenate, Bordeaux mixture alone, and lead arsenate alone in field 1, East Marion, Long Island, season of 1915 430
3. Curves showing relative wilt control of Bordeaux mixture and lead arsenate with date of first application as a variant in field 1, East Marion, Long Island, season of 1915 431

ALEYRODIDAE, OR WHITE FLIES, ATTACKING THE ORANGE, WITH DESCRIPTION OF THREE NEW SPECIES OF ECONOMIC IMPORTANCE

FIG. 1. *Aleurocanthus citriperdus:* *A*, Pupa case; *B*, egg; *C*, polygonal markings of egg; *D*, vasiform orifice of pupa case; *E*, spine from dorsum of pupa case; *F*, margin of pupa case; *G*, genitalia of adult male; *H*, forewing of male; *I*, antenna of pupa case; *J*, leg of pupa case; *K*, *L*, marginal teeth, much enlarged; *M*, central swollen spine from dorsal area 460
2. *Aleurocanthus woglumi:* *A*, Egg; *B*, polygonal markings of egg; *C*, pupa case; *D*, margin of pupa case; *E*, vasiform orifice of pupa case; *F*, forewing of adult female; *G*, same, showing variation in markings; *H*, costal margin at base of wing of female; *I*, forewing of male; *J*, male genitalia 462
3. *Aleurothrixus porteri*, *A. howardi*, and *A. floccosus:* *A*, *Aleurothrixus porteri:* Larva, first instar. *B*, *A. porteri:* Caudal spine of pupa case. *C*, *A. porteri:* Clasper of male. *D*, *A. porteri:* Egg. *E*, *A. howardi:* Caudal spine. *F*, *A. porteri:* Pupa case. *G*, *A. porteri:* Forewing of adult. *H*, *A. floccosus:* Vasiform orifice of pupa case. *I*, *A. porteri:* Vasiform orifice of pupa case. *J*, *A. howardi:* Vasiform orifice of pupa case. *K*, *A. porteri:* Margin of pupa case. *L*, *A. porteri:* Margin of early larva 467

RELATIVE WATER REQUIREMENT OF CORN AND THE SORGHUMS

FIG. 1. Curves of the evaporation at Garden City, Kans., for the growing period of 1915 474

CALIFORNIA GREEN LACEWING FLY

FIG. 1. The California green lacewing fly: Adult 515
2. *Chrysopa californica:* Eggs 518
3. *Chrysopa californica:* First instar 519
4. *Chrysopa californica:* Third instar 521
5. *Chrysopa californica:* Pupal case 522
6. *Chrysopa californica:* Pupa 522
7. *Chrysopa californica:* Pupa freshly emerged from its cocoon 523

STORAGE-ROTS OF ECONOMIC AROIDS

Page

FIG. 1. Spores of different storage-rot organisms: *A*, *Diplodia tubericola* from dasheen; *B*, *Diplodia tubericola* from sweet potato; *C*, *Diplodia gossypina* from cotton; *D*, *Diplodia* sp. from *Mangifera indica*: *E*, *Diplodia maclurae* from *Toxylon pomiferum*; *F*, *Fusarium solani*.. 552

LARVAL CHARACTERS AND DISTRIBUTION OF TWO SPECIES OF DIATRAEA

FIG. 1. *a*, Average angle formed by imaginary lines through bases of setæ of *Diatraea saccharalis crambidoides*; *b*, average angle formed by imaginary lines through bases of setæ of *D. zeacolella*..... 622

THE DISEASE OF POTATOES KNOWN AS "LEAK"

FIG. 1. Microscopical appearance of *Pythium debaryanum* isolated from potatoes affected with potato leak: *a*, Cell of a potato tuber showing fungus filaments therein; *b*, oogonia and antheridia; *c*, mycelium; *d*, germinating conidia..... 630

LIFE CYCLES OF THE BACTERIA

FIG. 1. Life cycle of *Bacillus azotobacter*. The broken straight lines divide the different types of growth indicated by the letters A to M. The Greek letters α to λ refer to subdivisions. The single and double pointed arrows indicate the development of one form from another. The four circles confine, in every case, all those forms which represent together a rather constant mode of life and which have been usually considered as bases for establishing separate species..... 678

MOTTLE-LEAF OF CITRUS TREES IN RELATION TO SOIL CONDITIONS

FIG. 1. Graphical presentation of the relationship between humus content of soil and percentage of mottled orange leaves..... 731

2. Graphical presentation of the relationship between the ratio of organic carbon to humus in the soil and the percentage of mottled orange leaves..... 734

3. Graphical presentation of the relationship between the ratio of humus to lime in the soil and the percentage of mottled leaves..... 735

4. Graphical presentation of the relationship between the mineral carbonates in the soil and the percentage of mottled lemon leaves..... 737

AGRICULTURAL VALUE OF IMPERMEABLE SEEDS

FIG. 1. Curves showing the rate of softening of impermeable red-clover seeds of different degrees of maturity..... 767

2. Curves showing the rate of softening and of germination of impermeable red-clover seeds of different lots..... 768

3. Curves showing the changes in the permeability of seeds in wet blotters and in dry storage for various periods..... 773

4. Curves showing the rate of softening of impermeable red-clover seeds under different temperature conditions..... 780

5. Curves of the seedling production in the field in 16 to 18 days and of the germination in chamber in 8 days..... 787

6. Curves of the seedling production in the field and of the germination in chamber in 12½ months..... 790

LIFE-HISTORY STUDIES OF CIRPHIS UNIPUNCTA, THE TRUE ARMY WORM

Page

FIG. 1. Diagram of relative amounts of foliage eaten in each larval instar by *Cirphis unipuncta*..... 801
2. Posterior extremity of male and female pupa: *a*, Male; *b*, female..... 812

THERSILOCHUS CONOTRACHELI, A PARASITE OF THE PLUM CURCULIO

FIG. 1. *Thersilochus conotracheli:* Cocoon..... 852
2. *Thersilochus conotracheli:* Egg..... 852
3. *Thersilochus conotracheli:* Larva of first instar..... 852
4. *Thersilochus conotracheli:* Ventral surface of head of first larval instar: *a*, Labrum; *b*, maxilla; *c*, labium; *d*, mandible..... 853
5. *Thersilochus conotracheli:* Full-grown larva of first instar..... 853
6. *Thersilochus conotracheli:* Larva of second instar..... 853
7. *Thersilochus conotracheli:* Full-grown larva of fifth instar..... 854
8. *Thersilochus conotracheli:* Face of full-grown larva: *a*, Labrum; *b*, mandible; *c*, maxilla; *d*, maxillary palpus; *e*, labium; *f*, labial palpus... 854
9. *Thersilochus conotracheli:* Pupa of female, and apex of abdomen of male pupa..... 854

EFFECT ON PLANT GROWTH OF SODIUM SALTS IN THE SOIL

FIG. 1. Diagram of the percentage of sodium carbonate added to the soil in experiment 1 (1913), with the percentage of carbonate and bicarbonate recovered and the total green weight of wheat obtained..... 860
2. Diagram of the percentage of sodium carbonate added to the soil in experiment 2 (1914), with the percentage of carbonate and bicarbonate recovered and the green weight of wheat obtained..... 862
3. Diagram of the percentage of sodium carbonate added to the loam soil in experiment 3 (1915), with the percentage of carbonate and bicarbonate recovered and green weight of wheat..... 864
4. Diagram of the percentage of sodium carbonate added to Monterey sand in experiment 3 (1915), with the percentage of carbonate and bicarbonate recovered and the total green weight of wheat obtained..... 864
5. Diagram of the percentage of sodium bicarbonate added to the soil in experiment 4 (1914), with carbonate and bicarbonate recovered, together with the total green weight of wheat obtained..... 865
6. Diagram of the decrease in growth of wheat seedlings in experiments 2 and 4 as affected by the total carbonate salts recoverable from the soil. 866
7. Diagram of the quantity of sodium chlorid added to the soil, with the quantity of chlorid recovered, and the total green weight of wheat obtained..... 867
8. Diagram of the quantity of sodium sulphate added to the soil in experiment 6, the quantity recovered, and the total green weight of wheat obtained..... 868

APHIDOLETES MERIDIONALIS, AN IMPORTANT DIPTEROUS ENEMY OF APHIDS

FIG. 1. *Aphidoletes meridionalis:* Eggs in situ on leaf of rape; *a*, egg..... 884
2. *Aphidoletes meridionalis:* Larva, dorsal view..... 886
3. *Aphidoletes meridionalis: a*, Cocoon formed on surface of ground; *b*, cocoon formed on a leaf blade..... 886
4. *Aphidoletes meridionalis:* Pupa, lateral view..... 887

INFLUENCE OF BARNYARD MANURE AND WATER UPON THE BACTERIAL ACTIVITIES OF THE SOIL

Page

FIG. 1. Curves of the ammonifying powers of soil in pots with varying quantities of manure and water 899
2. Curves of the nitrifying powers of soil in pots with varying quantities of manure and water 900
3. Curves of the number of colonies of bacteria developing from fallow soil with varying quantities of manure and water 904
4. Curves of the ammonifying powers of fallow soil with varying quantities of manure and water 907
5. Curves of the nitrifying powers of fallow soil with varying quantities of manure and water 910
6. Curves of the number of colonies of bacteria developing from cropped plats with varying quantities of manure 912
7. Curves of the ammonifying powers of soil of cropped plats with varying quantities of manure and water 915
8. Curves of the nitrifying powers of soil of cropped plats with varying quantities of manure and water 918
9. Diagram of the influence of manure on the yield and bacterial activities of a soil 920
10. Diagram of the influence of irrigation water on the yield and bacterial activities of a soil 921

PROGRESSIVE OXIDATION OF COLD-STORAGE BUTTER

FIG. 1. Diagram of gas apparatus used in the extraction and analysis of the air confined in butter 931

BACTERIOLOGICAL STUDIES OF A SOIL SUBJECTED TO DIFFERENT SYSTEMS OF CROPPING FOR 25 YEARS

FIG. 1. Curves of the nitrate formation in soil of fertility plats at Columbia, Mo., in 1913–14, after 28 days' incubation with the addition of 60 mgm. of nitrogen as cottonseed meal, but without the addition of calcium carbonate 962

STUDIES ON THE PHYSIOLOGY OF REPRODUCTION IN THE DOMESTIC FOWL.—XV. DWARF EGGS

FIG. 1. Diagram showing the percentage of the yearly total egg production (8-year average, 1899–1907) and the percentage of the total dwarf-egg production (8-year average, 1908–1916) which occurred during each month 1004
2. Diagram showing for the years 1911–12 and 1914–15 combined the percentage of the yearly total egg production and dwarf-egg production which occurred during each month and 100 times the percentage of the eggs produced each month which were dwarf 1005
3. Diagram showing the number of dwarf eggs which occurred in each tenth of a litter 1011
4. Diagram showing the number of dwarf eggs which occurred in each fifth of a litter 1014